

Effect of Polyploidy Induction on Biomass and Ginsenoside Accumulations in Adventitious Roots of Ginseng

Yun-Soo Kim¹, Eun-Joo Hahn², Hosakatte Niranjana Murthy^{2,3}, and Kee-Yoeup Paek^{2*}

¹Korea Ginseng Institute, Chung-Ang University, Ansong-shi 456-756, Korea

²Research Center for the Development of Advanced Horticultural Technology, Chungbuk National University, Cheongju 361-763, Korea

³Department of Botany, Karnatak University, Dharwad, 580 003, India

Adventitious roots of *Panax ginseng* C.A. Meyer (a natural tetraploid) were treated with 50 or 100 mg L⁻¹ colchicine for 12, 24, 36, 48, or 60 h to induce polyploid (octoploid) roots. The largest number of octoploid roots was obtained with a 100 mg L⁻¹ colchicine treatment over 60 h. To verify that ginsenoside was being accumulated in the developing tissues, the tetraploid (control) and octoploid roots were cultured for 40 d in Murashige and Skoog media that lacked NH₄NO₃ but was supplemented with 2 mg L⁻¹ naphthaleneacetic acid and 50 g L⁻¹ sucrose. Levels of fresh and dry biomass were greater in the octoploid roots. Although total ginsenoside and Rb-group ginsenoside contents were less in the octoploid roots than in the tetraploids, the former had a higher amount of Rg-group ginsenosides (especially Rg1). These results demonstrate the benefit that polyploid adventitious roots provide in enhancing the production of secondary metabolites in ginseng.

Keywords: colchicine, octoploid, *Panax ginseng* C.A. Meyer, tetraploid

Ginsenoside, a triterpenoid saponin, is a principal bioactive ingredient in *Panax ginseng*, a species with various physiological and pharmacological properties. The major types of ginsenosides are found in the Rb and Rg groups, which are derived from the 20(S) protopanaxadiol and 20(S) protopanaxatriol structures, respectively. Their main components include Rb1, Rb2, Rc, and Rd, from the Rb group; and Re, Rf, and Rg1 from the Rg group (Palazon et al., 2003).

As alternatives to field cultivation, ginseng cell and root cultures are used to produce ginsenosides because of their faster growth and stable metabolite production. However, ginsenoside contents in those cultured tissues are usually only about 30% of those derived from field-grown plants (Yu et al., 2002). Polyploidy is responsible for increases in cell size, a characteristic that leads to larger vegetative and reproductive organs. Polyploids are also responsible for improved secondary metabolite production (Dhawan and Lavania, 1996). Recently, this attribute has been induced in *Chamomilla recutita*, *Petunia*, and *Salvia miltiorrhiza*, where they are able to induce more flavonoid and terpenoid development than in their diploid counterparts (Gao et al., 1996; Griesbach and Kamo, 1996; Svehlikova and Repcak, 2000). There-

fore, the objective of this study was to use colchicine treatments to induce polyploidy (octoploidy) in the adventitious roots of *P. ginseng* in order to enhance root biomass and ginsenoside accumulation. We also compared root growth and ginsenoside contents between octoploid and naturally occurring tetraploid adventitious roots.

MATERIALS AND METHODS

Establishment of Adventitious Root Cultures and Induction of Polyploidy

Ginseng (*P. ginseng* C.A. Meyer) adventitious roots were induced and proliferated in an MS liquid medium (Murashige and Skoog, 1962) that lacked NH₄NO₃ but which was supplemented with 2 mg L⁻¹ naphthaleneacetic acid (NAA) and 50 g L⁻¹ sucrose (Kim et al., 2003). Cultures were maintained by shaking (100 rpm) at 25°C in the dark. To induce polyploidy (i.e., octoploidy), we inoculated 30 adventitious roots (3 cm long each) in 100-mL Erlenmeyer flasks containing 50 mL of culture media supplemented with 50 or 100 mg L⁻¹ of 0.2 µm filtered and sterilized colchicine (Sigma-Aldrich, USA). Each experiment lasted 12, 24, 36, 48, or 60 h. Following this colchicine treatment, the root tips were rinsed twice

*Corresponding author; fax +82-43-272-5369
e-mail paekky@chungbuk.ac.kr

tissues were treated with 50 mg L⁻¹ colchicine for 60 h. In contrast, 100 mg L⁻¹ colchicine induced chromosome doubling regardless of the treatment period, although the percent root survival decreased with increasing duration. For the 100 mg L⁻¹ colchicine treatment over 60 h, 73% of the roots showed doubling (Table 1).

Roots that had been determined to be octoploids (by chromosome count) were selected and sub-cultured in fresh culture media that lacked colchicine. Their ploidy level was confirmed by flow cytometry. The cells of barley, used as the internal standard, as well as those of the experimental materials were stained together and fed to the flow cytometer (Fig. 1). The internal standard (barley) showed two peaks (Peaks 1 and 3, Fig. 1A; Peak 1 and 2, Fig. 1B) whereas the tetraploid cells (control: natural ginseng roots) produced a histogram with only one peak (Peak 2, Fig. 1A). Compared with that tetraploid peak, the octoploid cells (from colchicine-treated roots) exhibited a peak twice the size of the channel position (Peak 3 in Fig. 1B), thereby confirming chromosome doubling as a result of colchicine application.

Growth and Ginsenoside Accumulation in Tetraploid and Octoploid Adventitious Roots

The octoploid adventitious roots were thick and bore fewer lateral branches (Fig. 2B), while the tetraploid adventitious roots were thin and had more laterals (Fig. 2A, Table 2). Fresh and dry biomasses of the octoploid adventitious roots were much greater (164.8 ± 11.5 FW and 13.8 ± 0.69 DW mg/root) compared with the naturally tetraploid roots (106.5 ± 3.9 FW and 11.7 ± 0.84 DW mg/root; Table 2). A similar increase in biomass has been reported in the polyploid (tetraploid) hairy roots of *Artemisia annua* when compared with its diploid roots (De Jesus-Gonzalez and Weathers, 2003).

Ginsenoside contents in both tetraploid and octoploid roots were analyzed by HPLC (Table 3). The amounts of total and Rb-group ginsenosides in the octoploid roots were lower (3.3 and 1.4 mg g⁻¹ DW,




Figure 1. Comparison of histograms obtained after flow cytometric analysis of tetraploid and octoploid adventitious roots of ginseng. **A**, Peaks 1 and 3 represent 2x and 4x nuclei of barley (internal standard); Peak 2 represents 4x nuclei of ginseng adventitious roots (tetraploid, without colchicine treatment). **B**, Peaks 1 and 2 represent 2x and 4x nuclei of barley (internal standard); Peak 3 represents 8x nuclei of ginseng adventitious roots (octoploid, after colchicine treatment).




Figure 2. Morphology of tetraploid (**A**) and octoploid (**B**) adventitious roots of ginseng, grown in MS medium (without NH₄NO₃) and supplemented with 2 mg L⁻¹ NAA and 50 g L⁻¹ sucrose.

Table 2. Growth and proliferation of tetraploid (4x) and octoploid (8x) adventitious roots after 40 d of culture.

Ploidy level	Fresh weight (mg)/ vessel	Dry weight (mg)/vessel	Dry weight/fresh weight (%)	No. lateral roots/root
4x ^z	106.5 ± 3.9 ^y	11.7 ± 0.84	10.9	11.9 ± 0.77
8x	164.8 ± 11.5	13.8 ± 0.69	8.4	4.4 ± 0.61

^zrepresents chromosome set per nucleus.

^yData are mean values ± SD from 30 adventitious roots.

Table 3. Ginsenoside accumulation in tetraploid (4x) and octoploid (8x) adventitious roots after 40 d of culture.

Ploidy level	Ginsenosides (mg g ⁻¹ dry wt)										Rb/Rg	
	Rb group				Rg group				TRb	TRg		Total
	Rb1	Rb2	Rc	Rd	Re	Rf	Rg1					
4x	0.6 ± 0.1	0.5 ± 0.1	0.4 ± 0.1	0.4 ± 0.0	0.6 ± 0.2	0.5 ± 0.0	0.7 ± 0.2	1.9 ± 0.2	1.8 ± 0.3	3.7 ± 0.3	1.1	
8x	0.4 ± 0.1	0.3 ± 0.2	0.3 ± 0.0	0.4 ± 0.1	0.5 ± 0.1	0.3 ± 0.1	1.1 ± 0.3	1.4 ± 0.2	1.9 ± 0.1	3.3 ± 0.2	0.7	

^yData are mean values ± SD from 3 replicates.

respectively) than in the tetraploid roots (3.7 and 1.7 mg g⁻¹ DW, respectively). However, Rg-group ginsenosides increased by at least 2%, with Rg1 ginsenoside, in particular, rising by 22% in the octoploid tissue. The overall ratio for the Rb:Rg group was 1.1 in octoploid roots and 0.7 in the tetraploids.

The various ginsenosides have different biological roles, e.g., the Rb group (protopanaxadiol saponins) shows anti-tumor activity (Zheng and Yang, 1994), while the Rg group (protopanaxatriol saponins) affects the motility of the intestine (Shin et al., 1997). Among the Rg-group ginsenosides, Rg1 (an active ingredient in ginseng), may have potential as a neuroprotective drug against Parkinson's disease (Chen et al., 2003) as well as being a hypertension, anti-fatigue medication. Therefore, the significantly enhanced level of Rg1 ginsenoside (up by 22%) in our octoploid roots demonstrates that this compound would be very advantageous for the intentional manipulation of the heterogeneity of ginsenosides in ginseng cell and root cultures. Quantitative differences in metabolites have been reported following polyploidy induction, for example, in the tetraploids of *Petunia*, *Salvia*, and *Artemisia*, which produce more secondary metabolites per gram of tissue than their diploid counterparts (Greisbach and Kamo, 1996; Svehlikova and Repcak, 2000; De Jesus-Gonzalez and Weathers, 2003). This enhanced accumulation of Rg1 ginsenoside in our octoploid adventitious roots of ginseng might also be exploited to develop clones for the production of higher amounts from the Rg1 group.

In conclusion, we have successfully induced octoploid ginseng adventitious roots via treatment with colchicine. To our knowledge, this is the first report on polyploid clones in that species. Here, the octoploid roots were responsible for higher accumulations of fresh and dry biomass. Although the total and Rb-group ginsenoside contents in the octoploid clones were lower than those measured in the tetraploids, we were able to obtain a greater amount of Rg-group ginsenosides, especially Rg1. These results indicate that induction of polyploidy, followed by screening for increased accumulations of individual ginsenosides,

could be beneficial to the production of secondary metabolites from ginseng cultures.

ACKNOWLEDGEMENTS

This work was supported by grants from the Korea Science and Engineering Foundation, Chungbuk National University, and the Rural Development Administration through the Biogreen 21 Project.

Received July 21, 2004; accepted October 22, 2004.

LITERATURE CITED

- Bulgakov VP, Lauve LS, Tchernoded GK, Khodakovskaya MV, Zhuravlev YN (2000) Chromosome variation in ginseng cells transformed with the *rolC* plant oncogene. *Russ J Genet* 36: 150-156
- Chen XC, Zhu YG, Zhu LA, Huang C, Chen Y, Chen LM, Fang F, Zhou YC, Zhao CH (2003) Ginsenoside Rg1 attenuates dopamine-induced apoptosis in PC12 cells by suppressing oxidative stress. *Eur J Pharmacol* 473: 1-7
- De Jesus-Gonzalez L, Weathers PJ (2003) Tetraploid *Artemisia annua* hairy roots produce more artemisinin than diploids. *Plant Cell Rep* 21: 809-813
- Dhawan OP, Lavania UC (1996) Enhancing the productivity of secondary metabolites via induced polyploidy: A review. *Euphytica* 87: 81-89
- Gao SL, Zhu DN, Cai ZH, Xu DR (1996) Autotetraploid plants from colchicine-treated bud culture of *Salvia miltiorrhiza* Bge. *Plant Cell Tiss Org Cult* 47: 73-77
- Griesbach RJ, Kamo KK (1996) The effect of induced polyploidy on the flavonoids of *Petunia* Mitchell. *Phytochem* 42: 361-363
- Grushvitskii IV (1961) *Zhenshen: Voprosy Biologii (Ginseng: Problems in Biology)*, Leningrad
- Kim YS, Hahn EJ, Yeung EC, Paek KY (2003) Lateral root development and saponin accumulation as affected by IBA or NAA in adventitious root cultures of *Panax ginseng* C. A. Meyer. *In Vitro Cell Dev Biol Plant* 39: 245-249
- Murashige T, Skoog F (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Plant Physiol* 15: 473-497

- Palazon J, Cusido RM, Bonfill M, Mallol A, Moyano E, Morales C, Pinol MT (2003) Elicitation of different *Panax ginseng* transformed root phenotypes for an improved ginsenoside production. *Plant Physiol Biochem* 41: 1019-1025
- Shin YH, Kim SC, Han JW, Oh JY, Jeong OM, Shin DH, Nah SY (1997) Ginseng protopanaxatriol saponins but not protopanaxadiol saponins inhibit spontaneous motility of intestine. *Koryo Insam Hakheochi* 21: 35-38
- Svehlikova V, Repcak M (2000) Variation of apigenin quantity in diploid and tetraploid *Chamomilla rectita* (L.) Rauschert. *Plant Biol* 2: 403-407
- Yu KW, Gao W, Hahn EJ, Paek KY (2002) Jasmonic acid improves ginsenoside accumulation in adventitious root culture of *Panax ginseng* C.A. Meyer. *Biochem Eng J* 11: 211-215
- Zheng GZ, Yang CR (1994) *Sanchi (Panax notoginseng) Biology and Application*. Science Press, Beijing